
 Europäisches Patentamt
 European Patent Office
 Office européen des brevets

Publication number:

0 210 761
A1

EUROPEAN PATENT APPLICATION

Application number: 86305125.6

Int. Cl. ⁴ C 12 N 9/02
A 61 K 37/50

Date of filing: 02.07.86

Priority: 05.07.85 JP 148922/85

Date of publication of application
04.02.87 Bulletin 87-6

Designated Contracting States
AT BE CH DE FR GB IT LI LU NL SE

Applicant: Takeda Chemical Industries, Ltd.
27, Doshomachi 2-chome Higashi-ku
Osaka-shi Osaka, 541(JP)

Inventor: Miyata, Kouichi
8-77, Seiwadainishi 1-chome
Kawanishi Hyogo 666-01(JP)


Inventor: Nakagawa, Yasushi
1-7, Midoridai 5-chome
Kawanishi Hyogo 666-01(JP)

Inventor: Nakamura, Masahira
408, Kitaimachi Kashiba-cho
Kitakatsuragi-gun Nara 639-02(JP)

Representative: Lewin, John Harvey et al,
Elkington and Fife High Holborn House 52/54 High
Holborn
London WC1V 6SH(GB)

Modified enzymes, production and use thereof.

The present invention provides a modified enzyme of the formula:



wherein SOD is a manganese-containing superoxide dismutase produced by a microorganism of the genus *Serratia*; n and m are the same or different and each is equal to about 7 to 700; R¹ and R² are the same or different and each is a hydroxy-protecting group; and x is equal to about 1 to 30.

The modified enzyme of the present invention shows increased antiinflammatory activity, is substantially free from immunogenicity and shows a prolonged half-life time and higher level in blood, hence it can advantageously be used as an antiinflammatory agent.


to about 1 to 30 and, preferably, about 5 to 15.

The polyethylene glycol used in accordance with the present invention should preferably have an average molecular weight of about 300 to about 30,000, more preferably about 1,000 to about 10,000.


As examples of the hydroxy-protecting groups represented by R^1 and R^2 , there may be mentioned alkyl groups containing 1-3 carbon atoms (e.g. methyl, ethyl, n-propyl, isopropyl), preferably methyl.

The modification of SOD with PEG may preferably be carried out via a triazine residue.


Such modification can be carried out by reaction of a mixture of a PEG of the general formula




wherein \bar{n} and R^1 are as defined above, and a PEG of the general formula



wherein \bar{m} and R^2 are as defined above, with a compound of the formula



and then reacting the resulting compound of the general formula



wherein \bar{n} , \bar{m} , R^1 and R^2 are as defined above, with SOD.

For producing the compound [IV], the reaction of

the mixture of compound [II] and compound [II'] with the compound [III] is preferably carried out under anhydrous conditions. Thus, the reaction is carried out in benzene or toluene at about 50° to about 80°C for about 5 15 to 50 hours.

The mixture of compound [II] and compound [II'] is preferably used in an amount such that the total number of moles of the compounds [II] and [II'] is twice the number of moles of the compound [III]. After the 10 reaction, aliphatic hydrocarbon-organic solvent, such as petroleum ether, is added to the reaction mixture to thereby cause precipitation of the compound [IV]. In some instances, gel filtration using an organic solvent (e.g. chloroform, acetonitrile, benzene, toluene), for 15 instance, may be employed combinedly.

For the production of the compound [I] by reaction of the compound [IV] with SOD, the reaction is carried out in an aqueous medium for about 2 to 24 hours under mild conditions to avoid enzyme inactivation, for 20 example at room temperature or below and at a pH of about 8 to about 10. After reaction, the reaction mixture is subjected to gel filtration and ultra-filtration in an aqueous solution to separate the compound [I] and the remaining compound [IV].

25 By these procedures, the compound [I], a PEG-modified SOD, is obtained, hence hereinafter sometimes referred to as "PEG-SOD".

The compound [I] according to the invention has sufficiently high SOD activity. Testing of the compound 30 for evaluating antiinflammatory agents [I] by such model experiments as the carrageenin abscess model, delayed type IV allergic reaction with ovalbumin, Masugi nephritis model and radiation damage has revealed that the PEG-modified SOD is generally superior in pharmacological effects to the original SOD. 35

As compared with the original SOD, the compound [I] has the following advantageous features:

- (1) It increases antiinflammatory activity.
- (2) It decreases immunogenicity.
- 5 (3) Its half-life time in blood circulation is longer.
- (4) One time injection of it induces immunological tolerance.
- (5) Blood level of SOD is much higher after intraduodenal administration.

10 Furthermore, the compound [I] has low toxicity.

In this manner, the compound [I] has excellent activity and properties and, in addition, is low in toxicity. Therefore, it can be used advantageously as an antiinflammatory agent.

15 In using the compound [I] according to the invention as an antiinflammatory agent, it is administered either orally or parenterally for the purpose of treating various acute or subacute inflammations (e.g. edema, (e.g. arthritis, bronchitis, bruise-caused edema, burn, 20 radiation)) in mammals (e.g. mouse, rat, rabbit, cat, dog, monkey, human), for instance.

For administration, the compound [I] is mixed with a pharmacologically acceptable carrier, excipient, diluent, etc., and made up into preparations for oral 25 administration, such as tablets or capsules, or into preparations for parenteral administration, such as injections, and is administered to the above mammals in such preparation form by the per se known method.

The daily dose of the compound [I] is such that 30 the enzyme quantity in the compound [I] amounts to about 0.1 to about 100 mg, preferably about 0.5 to 50 mg, per kilogram of body weight.

Brief Description of the Drawings

Fig. 1 shows the results of immunological tolerance 35 induction as obtained in Test Example 6.

Fig. 2 shows the results of plasma level of SOD activity after intraduodenal administration to rats as disclosed in Test Example 7.

Examples

5 The following working examples, reference example and test examples are further illustrative of the present invention.

Example 1 Production of PEG-SOD

10 Cyanuric chloride (730 mg) was added to a mixture of 40 g of monomethoxy polyethylene glycol (average molecular weight: 5,000), 200 ml of benzene, 20 g of anhydrous sodium carbonate and 10 g of molecular sieve 3A (Wako Pure Chemical Industries, Japan), and the reaction was carried out by stirring at 80°C for 20
15 hours. Thereafter, 400 ml of petroleum ether was added to thereby cause precipitation of the activated PEG. The precipitate was dissolved in benzene, followed by the precipitation by the same procedure. After three-time repetitions of this precipitation procedure, the
20 precipitate (activated PEG) was dried in a desiccator under reduced pressure.

 A 1.5 g portion of the above activated PEG was added to 15 mg of an SOD prepared from Serratia marcescens ATCC 21074 by the procedure described in EP
25 Publication (laid open) No. 70656 (The mole ratio of the activated PEG to the SOD was 300.) The reaction was carried out at 4°C for 2 hours in 5.0 ml of 0.1 M borate buffer (pH 9.0) and then terminated by adding 30 ml of 0.1 M phosphate buffer (pH 7.0). The residual PEG was
30 removed by ultrafiltration [Amicon (USA) membrane YM-10]. The concentrate (2.0 ml) of PEG-SOD was purified by gel filtration using a Sephacryl S-200 (Pharmacia, Sweden) column (5 x 73 cm). The thus-obtained PEG-SOD, weighed about 15 mg, was modified
35 7-12 amino residues of the original SOD, and its SOD

activity retained about 50%. The ultraviolet absorption at 280 nm was increased by less than 5% of the original. At shorter wavelengths, the absorptions due to the triazine residue were observed.

5 Example 2 Production of PEG-SOD

PEG-SODs were prepared by the same method as in Example 1 using monomethoxy polyethylene glycols differing in molecular weight (average molecular weight: 350, 750 and 1,900, respectively) with the SOD used in
10 Example 1. The results thus obtained are shown in Table 1.

Table 1

	Specific activity (units/mg protein)	Amino group modification percentage
15 SOD (original)	100%	0%
PEG (350)-SOD	39%	48%
PEG (750)-SOD	41%	48%
PEG (1,900)-SOD	41%	48%

20

Note: The value in the parentheses after "PEG" indicates the average molecular weight of the PEG used.

Reference Example 1 Preparation of dextran-SOD

25 In 50 ml of distilled water was dissolved 1 g of dextran (average molecular weight: 60,000-90,000), followed by addition of about 0.3 g of cyanogen bromide. The reaction was carried out at 20°C for 30 minutes while the pH was maintained at 10.5 by addition of 5 N
30 NaOH. Then, the pH was adjusted to 9.0 with NaHCO₃, the SOD used in Example 1 (114 mg in 13 ml of distilled water) was added, and the reaction was carried out furthermore at 4°C for 20 hours. The reaction mixture was concentrated to 3 ml by ultrafiltration (using
35 YM-10) and the remaining SOD was removed by gel

filtration using a Sephacryl S-200 column (5 x 73 cm). Thus was obtained about 0.7 g of dextran-SOD. The dextran-SOD modified with 34% of the amino groups had a residual enzymatic activity of 67% of the original SOD.

5 Test Example 1 Antiinflammatory activity

The carrageenin abscess test was used as an evaluation method of antiinflammatory activity. Thus, as described in *Acta Pharmacologica Japonica*, 84, 337 (1984), a 2% (w/v) carrageenin solution was subcutaneously injected into the dorsal region of rats. Thirty minutes later, the enzymes show in Table 2 were each administered intravenously. After further 24 hours, each abscess was excised and weighed for comparison with the control group (given physiological saline).

In this and the subsequent test examples, the SOD used was same used in Example 1 and the dextran-SOD used was the product obtained in Reference Example 1 described above.

20 Table 2 Antiinflammatory activity
(carrageenin abscess method)

	Dose ¹⁾	Number of animal	Abscess weight ²⁾	Inhibition
25 Control	-	10	2.117 ± 0.11 g	-
SOD	2 mg/kg	10	2.065 ± 0.13 g	2.5%
Dextran-SOD	2 mg/kg	10	1.695 ± 0.12 g ³⁾	20.5%
30 PEG-SOD ⁴⁾	2 mg/kg	10	1.701 ± 0.11 g ³⁾	19.7%

Notes: 1) Expressed in terms of the number, the doses were same as SOD activity.

2) Mean ± standard error

35 3) P < 0.05 (Student's T test)

4) The PEG-SOD obtained in Example 1 was used.

As shown in Table 2, the modified enzymes, dextran-SOD and PEG-SOD, increased antiinflammatory activity as compared with the original SOD and the efficacy of the modified enzymes showed statistically significant as compared with that of the control group (given physiological saline).

Test Example 2 Immunogenicity

The method described in Journal of Immunological Methods, 14, 381 (1977) was used. Each enzyme (20 μ g as SOD) was emulsified with Freund's complete adjuvant (FAC) and the emulsion (0.2 ml) was administered intraperitoneally to A/J mice, followed by additional administration on days 14 and 28. Blood samples were taken from the mouse orbital vein every 7 day interval after the day of the first administration. The sera from the blood samples were evaluated for antibody production by the passive cutaneous anaphylaxis (PCA) reaction method. Rats were intradermally injected with a diluted serum (0.1 ml), followed by injecting 2 ml of a solution containing 0.5 mg of SOD and 20 mg of Evans blue 4 hours later and the vascular permeability visualized by the dye was determined. The results thus obtained are shown in Table 3. The values in the table indicate the maximum serum dilution factor capable of giving positive PCA reaction.

Table 3 Immunogenicity testing

	Number of animal	PCA titer judged on				
		Day 7	Day 14	Day 21	Day 28	Day 35
SOD	8	8	32	256	256	256
Dextran-SOD	8	0	256	128	128	256
PEG-SOD ¹⁾	8	0	0	0	0	0

- 10 -

Note 1) The PEG-SOD used was the product obtained in Example 1.

As shown in Table 3, the PEG-SOD showed no antibody production while the original SOD and dextran-SOD caused antibody production against SOD. In a test conducted by the similar method, no antibody production against PEG-SOD was observed in the PEG-SOD administered rats.

Test Example 3 Half-life time in blood/circulation
Each enzyme (5 mg/kg as SOD) was intravenously injected (1 ml) into rats (3 per group). Blood samples were taken from the caudal vein at time intervals and measured for serum SOD activity, and the half-life time was determined based on the first order reaction rate constant.

Table 4 Half-life time in blood circulation

SOD	1.4 hours
PEG-SOD ¹⁾	14.2 hours

Note 1) The PEG-SOD obtained in Example 1 was used.

As shown in Table 4, the half-life time of PEG-SOD in the blood circulation was about 10 times longer as compared with that of the original SOD.

Test Example 4 Half-life time in blood circulation

The half-life time of SOD and the PEG-SODs obtained in Examples 1 and 2 were determined by the same described in Test Example 3. The results are shown in Table 5.

- 11 -

Table 5 Half-life time in blood circulation

		Half-life time ¹⁾
5	SOD	1.4 hours
	PEG(350)-SOD ²⁾	6.3 hours
	PEG(750)-SOD ²⁾	8.7 hours
	PEG(1,900)-SOD ²⁾	11.2 hours
10	Notes: 1) Determined by the same method described in Test Example 3.	
	2) The PEG-SODs obtained in Example 2 were used.	

As shown in Table 5, the SOD's modified with various PEG's showed a prolonged half-life time, compared with that of the original SOD.

Test Example 5 Immunogenicity

The PEG-SODs obtained in Example 2 were tested for immunogenicity by the same method described in Test Example 2. The results are shown in Table 6.

Table 6 Immunogenicity test

		PCA titer judged on				
		Number of animal	Day 7	Day 14	Day 21	Day 28
25	SOD	8	4	128	128	512
	PEG(350)-SOD ¹⁾	8	0	8	32	256
	PEG(750)-SOD ¹⁾	8	0	4	16	32
	PEG(1,900)-SOD ¹⁾	8	0	0	0	0

Note 1) The PEG-SODs obtained in Example 2 were used.

As is evident from Table 6, the modification of SOD with PEG reduced antibody production. Especially for the PEG(1,900)-SOD, no antibody production was noted.

Test Example 6 Induction of immunological tolerance

A/J mice (8 per group) were intravenously injected with the PEG-SOD obtained in Example 1 at a dose of 20 μ g SOD/mouse. An emulsion of 20 μ g of SOD with Freund's complete adjuvant (FCA)/mouse was intraperitoneally administered 1, 3 and 5 weeks later. The titers of anti-SOD IgE antibody were determined by the PCA reaction described in Test Example 2. The results are shown in Fig. 1. As is apparent from Fig. 1, the initial administration of the PEG-SOD (shown by ---o---) completely suppressed the IgE antibody production against SOD even after the subsequent administrations of SOD. (Note: IgE = Immunoglobulin E.)

For the control, 20 μ g of SOD/mouse was intravenously injected into A/J mice (8 per group) and an emulsion of 20 μ g of SOD in FCA was administered intraperitoneally 1, 3 and 5 weeks later. The results are also shown in Fig. 1 by --- Δ ---. As is apparent from Fig. 1, the initial administration of the original SOD elicited the IgE antibody production against SOD. The data indicated by --- \bullet --- in Fig. 1 show the pattern of IgE antibody production against SOD when physiological saline was intravenously injected into A/J mice (8 per group) followed by intraperitoneal administration of an emulsion of 20 μ g of SOD in FCA 1, 3 and 5 weeks later.

As is evident from the above results, the PEG-SOD induced immunological tolerance more strongly than the original SOD.

Test Example 7 Plasma level of SOD activity after intraduodenal administration

PEG-SOD obtained in Example 1 and SOD were administered intraduodenally through a cannula to rats (male, Jcl:Sprague-Dawley, 7 week old) at a dose of 100 mg as SOD/kg. Blood was taken from the rat tail vein 0,

1, 2, 4, 6, 10 and 24 hours after the administration and SOD activity in the plasma was assayed. The results are shown in Fig. 2. The plasma SOD level of the PEG-SOD administered rats presented by —○— was much
5 higher than that of the SOD administered rats presented by ---Δ--- throughout the time course. Each point represents the mean with standard error (SE) of 5 rats in PEG-SOD and 3 rats in SOD.

10

15

20


25

30

35

What is claimed is:

1. A modified enzyme of the formula:



wherein SOD is a manganese-containing superoxide dismutase produced by a microorganism of the genus Serratia; \bar{n} and \bar{m} are the same or different and each is equal to about 7 to 700; R^1 and R^2 are the same or different and each is a hydroxy-protecting group; and \bar{x} is equal to about 1 to 30.

2. The enzyme according to Claim 1, wherein \bar{n} and \bar{m} are the same.


3. The enzyme according to Claim 1, wherein \bar{n} and \bar{m} each is equal to about 7 to 250.

4. The enzyme according to Claim 1, wherein R^1 and R^2 are the same.

5. The enzyme according to Claim 1, wherein R^1 and R^2 each is C_{1-3} alkyl.

6. The enzyme according to Claim 1, wherein SOD is produced by Serratia marcescens ATCC 21074.

7. A method of producing the enzyme according to Claim 1, which comprises reacting SOD with the compound of the formula:



wherein \bar{n} , \bar{m} , R^1 and R^2 are defined in Claim 1.

8. A pharmaceutical composition which contains the enzyme according to Claim 1 and a pharmacologically acceptable carrier, excipient or diluent therefor.


9. The composition according to Claim 8, which is formulated to a tablet or capsule for oral administration.

10. A method for treatment of inflammation in a mammal, which comprises administering an effective amount of the enzyme according to Claim 1 to the mammal.

1/2

0210761


Fig. 1



2/2

0210761

Fig. 2





European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT
which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

0210761

Application number
EP 86 30 5125

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X	RADIATION RESEARCH, volume 93, 1983, pages 581-587 B.H. GRAY: "Radioprotection by polyethylene glycol-protein complexes in mice" * page 582 * --	1-7	C 12 N 9/02 A 61 K 37/50
X	CHEMICAL ABSTRACTS, volume 93, no. 15, October 13, 1980 page 64, abstract 143000f COLUMBUS, OHIO (US) P.S. PYATAK et al.: "Preparation of a polyethylene glycol: superoxide dismutase adduct, and an examination of its blood circulating life and anti-inflammatory activity" & Res. Commun. Chem. Pathol. Pharmacol. 1980, 29(1), 113-27 * abstract * --	1-9	TECHNICAL FIELDS SEARCHED (Int. Cl. 4) C 12 N 9 A 61 K 37
INCOMPLETE SEARCH		./.	
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-9 Claims searched incompletely: 10 Claims not searched: 10 Reason for the limitation of the search:</p> <p>Method for treatment of the human or animal body by surgery or therapy (see art. 52 (4) of the European Patent Convention).</p>			
Place of search THE HAGUE		Date of completion of the search 25-09-1986	Examiner SKELLY
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

0210761

Application number
EP 86 30 5125 -2-

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Category	Citation of document with indication where appropriate, of relevant passages	Relevant to claim	
X	CHEMICAL ABSTRACTS, volume 94, no. 7, February 16, 1981 page 226, abstract 43310j COLUMBUS, OHIO (US) F.F. DAVIS et al.: "Soluble, nonantigenic polyethylene glycol-bound enzymes" & Polym. Prepr. Am. Chem. Soc. Div. Polym. Chem. 1979, 20(1) 357-60 (Eng). * abstract *	1-9	
Y	CHEMISTRY LETTERS, no. 7, July 1980, pages 773-776 The Chemical Soc. of Japan (JP) A. MATSUSHIMA et al.: "Modification of E.Coli asparaginase with 2,4-bis(0-methoxypolyethylene glycol)-6-chloro-S-triazine (activated Peg ₂); disappearance of binding ability towards anti-serum and retention of enzymic activity" * whole document *	7	TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
Y, D	EP - A - 0 045 222 (TAKEDA CHEMICAL INDUSTRIES) * whole document *	7	
A, D	EP - A - 0 070 656 (TAKEDA CHEMICAL INDUSTRIES) * whole document *	1-9	
A	LA RECHERCHE, volume 10, no. 106 December 1979, page 1269 PARIS (FR) M. MICHELSON: "Une enzyme qui nous veut du bien" * whole document *	1-9	